

THE 2MASS WIDE-FIELD T DWARF SEARCH. I. DISCOVERY OF A BRIGHT T DWARF WITHIN 10 PC OF THE SUN

ADAM J. BURGASSER^{1,2}, J. DAVY KIRKPATRICK³, MICHAEL W. McELWAIN¹, ROC M. CUTRI³,
ALBERT J. BURGASSER⁴, & MICHAEL F. SKRUTSKIE⁵

Accepted by AJ for February 2003

ABSTRACT

We present the discovery of a bright ($J = 13.94 \pm 0.03$) T dwarf, 2MASS 1503+2525, identified in a new, wide-field search for T dwarfs using the recently completed Two Micron All Sky Survey (2MASS). The 1–2.5 μm spectrum of this object exhibits the strong H_2O and CH_4 bands characteristic of mid- and late-type T dwarfs, and we derive a spectral type of T5.5 using both the Burgasser et al. and the Geballe et al. classification schemes. Based on its spectral type and the absolute magnitudes of known T dwarfs, we estimate the distance of this object as 8 ± 3 pc if it is single, likely within 10 pc of the Sun. Our new 2MASS search, which covers 74% of the sky and greatly expands on earlier color constraints, should identify 15–25 new T dwarfs with $J \leq 16$. Combined with the 20 known members of this class that already fall within our search criteria, our new sample will provide improved statistics for such key quantities as the binary fraction and the field substellar mass function. Furthermore, multiple detections from overlapping 2MASS scans provide multiple epoch astrometry and photometry, and we present proper motions for five T dwarfs in our sample.

Subject headings: Galaxy: solar neighborhood — infrared: stars — stars: individual (2MASS J15031961+2525196) — stars: low mass, brown dwarfs

1. INTRODUCTION

T dwarfs are low-temperature ($T_{\text{eff}} \lesssim 1300\text{--}1500$ K) brown dwarfs exhibiting distinct absorption bands of CH_4 in the near-infrared H- and K-bands (Kirkpatrick et al. 1999; Burgasser et al. 2002b; Geballe et al. 2002). They are distinguished from warmer L-type (Kirkpatrick et al. 1999; Martín et al. 1999) and M-type dwarfs in the near-infrared by the presence of these molecular features, in addition to significant H_2O and collision-induced (CIA) H_2 absorption (Saumon et al. 1994). Since the discovery of the prototype of this class, Gliese 229B (Nakajima et al. 1995; Oppenheimer et al. 1995), over 30 T dwarfs have been identified, most in the wide-field Two Micron All-Sky Survey (Skrutskie et al. 1997, hereafter 2MASS) and Sloan Digital Sky Survey (York et al. 2000, hereafter SDSS). Classification schemes for these objects based on near-infrared spectral morphology have been defined by Burgasser et al. (2002b) and Geballe et al. (2002).

T dwarfs identified in wide-field surveys are all likely within 25 pc of the Sun, the limiting distance for the NASA/NSF NStars Project. Indeed, brown dwarfs in general may constitute a significant population in the Solar Neighborhood, roughly equal in number to the stellar population (Reid et al. 1999; Chabrier 2002). They have generally eluded detection, however, because of their intrinsic faintness. As the Solar Neighborhood is the basis for

many studies of stellar atmospheres, star formation and evolution, initial mass function, and Galactic structure, characterizing the local substellar population is of some importance. This is especially true at closer distances, such as the 10 pc RECONS sample (Henry et al. 1997), where the stellar census is far more complete, but few L- or T-type stars and brown dwarfs are currently known.

In this article, we present the discovery of a bright T dwarf, 2MASS J15031961+2525196⁶ (hereafter 2MASS 1503+2525), identified in a new search of the all-sky 2MASS Working Point Source Database (WPSD). The spectrophotometric distance of this object places it within 10 pc of the Sun. In §2 we describe our new wide-field T dwarf search and compare it to previous search efforts. In §3 we discuss the identification of 2MASS 1503+2525 in our search and its subsequent confirmation as a T dwarf using the Lick 3m Gemini Infrared Camera. In §4 we derive the spectral type and spectrophotometric distance of 2MASS 1503+2525. We discuss our results in §5.

2. THE 2MASS WIDE-FIELD T DWARF SEARCH

Our current program is a continuation of a previous 2MASS search (Burgasser et al. 2002b) conducted when this survey was roughly 50% complete. 2MASS is ideal for identifying T dwarfs, as the JHK_s filters it has employed sample the peak of the spectral energy distribution of cool

¹ UCLA Division of Astronomy & Astrophysics, 8965 Math Science Bldg., 405 Hilgard Ave., Los Angeles, CA, 90095-1562; adam@astro.ucla.edu, mcelwain@astro.ucla.edu

² Hubble Fellow

³ Infrared Processing and Analysis Center, M/S 100-22, California Institute of Technology, Pasadena, CA 91125; davy@ipac.caltech.edu, roc@ipac.caltech.edu

⁴ John J. McCarthy Observatory, P.O. Box 1144, New Milford, CT 06776; aburgasser@mccarthyobservatory.org

⁵ University of Virginia, Department of Astronomy, P.O. Box 3818, Charlottesville, VA 22903; skrutskie@virginia.edu

⁶ Throughout this paper, we cite designations as they appear in the final 2MASS Point Source Catalog (PSC), given as “2MASS Jhhmmss[.ss±ddmmss[.s]”. The suffix conforms to IAU nomenclature convention and is the sexagesimal Right Ascension and declination at J2000 equinox.

brown dwarfs, enabling us to probe relatively deeply and over a large area of sky. A total of 17 T dwarfs were detected in our earlier search in three separate samples: a primary sample covering $16,620 \text{ deg}^2$, requiring $J < 16$, $J-H < 0.3$, and $H-K_s < 0.3$ for candidates selected from the 2MASS WPSD; and two auxiliary samples selected from 2MASS Incremental Release Data with smaller search areas but somewhat different color criteria. However, the overall stringent color cuts used in this search prevented our detecting early- and mid-type T dwarfs (Leggett et al. 2000), while our small sample resulted in large statistical uncertainties for population statistics such as the T dwarf space density and substellar mass function (Burgasser 2001; Burgasser et al.; in preparation).

Our current search effort, using the 2MASS WPSD, attempts to address these shortcomings in two ways. First, we have expanded our color criteria to $J-H \leq 0.3$ or $H-K_s \leq 0$, allowing detection of objects with redder near-infrared colors (and hence earlier spectral type), or faint objects with larger photometric uncertainties that were not identified in our first primary sample. These color constraints are required to exclude an overwhelming number of background stars, as discussed below, and effectively limit our search to spectral types roughly T4 and later (Burgasser et al. 2002b; Leggett et al. 2002). We continue to impose signal-to-noise (S/N), photometric completeness, and reliability constraints by selecting only those candidates with $J \leq 16$ ($S/N \sim 10$ completeness limit) and both J- and H-band detections. Second, with the recent completion of 2MASS survey operations, we are now able to search the entire sky for T dwarfs, excluding only those regions with very high source densities and hence likelihood of source confusion: the Galactic Plane ($|b| < 15^\circ$), the Large and Small Magellanic Clouds, and various dense or obscured stellar regions such as 47 Tuc and the Orion Nebular Cluster. We have also excluded the Equatorial poles ($|\delta| > 88^\circ$) because of restrictions on observational followup. In total, our current search encompasses $30,400 \text{ deg}^2$, or 74% of the sky, increasing our previous sample area by over 80%. We expect to find roughly 15 new T dwarfs in this sample based on the increased areal coverage alone, and up to 10 additional red and/or faint T dwarfs based on the number of objects found in the Burgasser et al. (2002b) auxiliary samples.

In Figure 1 we plot the $J-H$ and $H-K_s$ Hess diagram (Hess 1924) of our 264,068 2MASS-selected candidates, selected from over 1.3 billion sources in the 2MASS WPSD, along with the 2MASS colors of T dwarfs detected by 2MASS and SDSS (Table 7 in Burgasser et al., 2002; and Table 4 in Geballe et al., 2002; all photometry from 2MASS). The density of candidates is highest near our color constraints, particularly at the early-type end of the Bessell & Brett (1988) dwarf track and near the turnoff between the dwarf (late-K to early-M type) and giant tracks. The vast majority of these candidates are likely faint background stars, which are densely centered at and

around the Bessell & Brett (1988) tracks. While our 2MASS-selected candidates are required to have no optical counterparts within $5''$ in the USNO A2.0 catalog (Monet et al. 1998), based on the observed extreme red colors of known T dwarfs ($R-J \gtrsim 9$; Golimowski et al. 1998), a substantial number of faint sources, proper motion stars, closely-separated visual binaries, and optically red sources (the USNO A2.0 catalog requires detections in both R and B bands) remain. Hence, we visually examine multiple-epoch Digital Sky Survey (DSS) images of all candidate fields to exclude these contaminants. Of the roughly 70,000 candidates so far examined, 99.5% have been eliminated in this manner. The remaining steps of our search process, as discussed in Burgasser et al. (2002b), include follow-up near-infrared imaging for the elimination of minor planet contaminants, optical imaging to exclude faint background stars, and near-infrared spectroscopy for final identification. These follow-up observations are currently underway.

Our color and magnitude constraints encompass 20 T dwarfs previously identified by 2MASS and SDSS, listed in Table 1⁷. This sample is already larger than that studied by Burgasser et al. (2002b). Furthermore, seven of these sources have multiple detections in the 2MASS WPSD due to overlapping or repeated scans, as listed in Table 2. These observations, separated by as much as 2 years, enable us to derive mean proper motions for six of the T dwarfs⁸, assuming positional uncertainties of $0''.3$ ⁹. Note that these values have not been corrected for Solar motion. Four of the objects, 2MASS 0937+2937, SDSS 1346-0031, Gliese 570D, and 2MASS 2339-1352, have motions of at least $0''.5 \text{ yr}^{-1}$. The substantial motions of these objects are not unexpected, given that T dwarfs detected by 2MASS are all likely within 20 pc of the Sun.

A comparison of the multiple-epoch photometry for the objects listed in Table 2 gives some hint of photometric variability. Reasonably significant (maximum $\Delta M \gtrsim 2\sigma$) variations are seen at J-band for SDSS 0926+5847 (2.8σ), 2MASS 1237+6526 (2.1σ), Gliese 570D (2.8σ), and 2MASS 2339+1352 (3.6σ); and at H-band for SDSS 1346-0031 (2.3σ). It is possible that these photometric variations are intrinsic, as has now been observed for a number of brown dwarfs (Bailer-Jones & Mundt 1999, 2001; Clarke, Tinney, & Covey 2002; Gelino et al. 2002, Enoch, Brown, & Burgasser, in preparation). Burgasser et al. (2002a) have hypothesized that these variations arise from the motions and evolution of patchy condensate clouds in the upper atmospheres of early- and mid-type T dwarfs; indeed, the variations appear to be greatest at J-band, as predicted, and possibly more pronounced in the T4.5 SDSS 0924+5847 and T5.5 2MASS 2339+1352 (note the equally strong variation in the T8 Gliese 570D, however.) Such low-level photometric variability as measured by the 2MASS J-band filter must be considered carefully, however, as the filter bandpass extends well into the $1.4 \mu\text{m}$ telluric H_2O band. This results in substantial variations in

⁷ Gliese 229B is not listed in Table 1, for at a separation of $7''.8$ Nakajima et al. (1995) it is obscured by its brighter primary in the 2MASS imaging data. Distant T dwarfs, such as NTTDF 1205-0744 (Cuby et al. 1999), IFA 0231-Z1 (Liu et al. 2002), and SOri 0538-0236 (Zapatero Osorio et al. 2002) are too faint to satisfy our magnitude constraints.

⁸ The proper motion for Gliese 570D has been previously reported in Burgasser et al. (2000a), confirming that object's association with Gliese 570ABC.

⁹ Characterization of the astrometric and photometric properties of 2MASS point sources can be found in Cutri et al. (2002); <http://www.ipac.caltech.edu/2mass/releases/second/doc/explsup.html>.

the J-band zeropoint (up to 0.1 mag) from night-to-night and within a single night (Cutri et al. 2002). The source of these fluctuations, likely changes in observing conditions (terrestrial weather and water vapor), can potentially affect the photometric calibration of late-L and T dwarfs, as their spectral energy distributions show far more structure in the J-band than earlier-type calibration stars. Such second-order extinction effects could therefore be responsible for the apparent variability seen, and a more directed program is required to validate these observations. Regardless, our complete sample will provide time-resolved astrometric and potentially photometric data for a few individual T dwarfs, in addition to improved population statistics.

3. IDENTIFICATION OF 2MASS 1503+2525

3.1. Initial Selection

2MASS 1503+2525 is one of the first objects examined in our new wide-field search campaign. Astrometry and photometry for this object are listed in Table 1, while Figure 2 shows the 2MASS 1503+2525 field as imaged by POSS-I (Wilson 1952), POSS-II (Reid et al. 1991), and 2MASS. This object is one of our brightest candidates, just fainter than the brightest known T dwarf 2MASS 0559–1404 (Burgasser et al. 2000b). The absence of an optical source at the position of 2MASS 1503+2525 implies $R-J \gtrsim 6$ (Reid et al. 1991), while its near-infrared colors ($J-H = 0.08 \pm 0.05$, $H-K_s = -0.11 \pm 0.08$) are similar to the T6.5 Gliese 229B (Leggett et al. 1999). The optical data also rule out the possibility that this candidate is an earlier-type high-velocity proper motion star, as no such counterpart is seen at either epoch. Note that the bright star $18''$ North and $103''$ East of 2MASS 1503+2525 is the M5 variable star EX Boo (a.k.a. IRAS 15012+2537, BD+25 2864, HIP 73662), classified as a normal giant/supergiant by Guglielmo et al. (1997). A lower limit on this object’s Hipparcos distance, $d > 680$ pc (Perryman et al. 1997), eliminates the possibility that it is associated with the T dwarf.

3.2. Lick Gemini Infrared Camera Observations

We imaged 2MASS 1503+2525 in the near-infrared on 27 June 2002 (UT) using the Gemini Twin-Arrays Infrared Camera (McLean et al. 1993), mounted on the Lick Observatory Shane 3m Telescope. Light cirrus was present and seeing was $1''.2$. Short integration (20 sec), J- and K-band images unambiguously confirmed the presence of 2MASS 1503+2525 at its 2MASS position, and no secondary component was seen at the seeing spatial resolution of the data.

We subsequently obtained low-resolution ($\lambda/\Delta\lambda \sim 500$) JHK spectra using the Gemini Camera’s grisms on two separate nights, 28 June 2002 and 19 August 2002 (UT). Conditions for the first night (J- and K-band spectra) were affected by heavy cirrus and $2''$ seeing, while smoke and light cirrus ($1''.5$ seeing) were present on the second night (H-band spectrum). For both sets of observations, 2MASS 1503+2525 was first acquired in imaging mode and placed into a $1''.4$ slit. Sets of noddred ($30''$) pairs were obtained with individual integration times of 180–300 sec, for total integrations of 1800 sec at J- and K-bands, and 2000 sec at H-band. Nod pair sets of the F8 V HD 133460 were

obtained for calibration immediately after the target observations and at similar airmass.

Spectral images were pairwise subtracted; divided by a normalized, median-combined set of flat-field quartz lamp observations; and corrected for bad pixels by linear interpolation. Spatial curvature was measured using the calibrator star dispersion lines, and spectral curvature was measured from Argon lamp spectra obtained during each of the observing runs. These lamp spectra were also used for wavelength calibration, using the identifications of Stringov & Sventitskii (1968). Target and calibrator spectra at each nod position were then individually extracted and interpolated onto a common wavelength scale. Relative flux calibrations and telluric corrections were made by dividing target spectra by calibrator spectra at each nod position, interpolating over Paschen and Brackett Hydrogen lines. The resulting individual spectra for each band were multiplied by a 6095 K blackbody (Tokunaga 2000, corresponding to spectral type F8 V) and averaged. 2MASS magnitudes were used to flux calibrate the separate bands by the prescription of Burgasser et al. (2002b).

Figure 3 plots the reduced Gemini Camera grism spectrum of 2MASS 1503+2525. Signal-to-noise ratios are roughly 17 at J, 13 at H, and 9 at K. CH_4 bands at 1.15, 1.4, and $2.2 \mu\text{m}$, and H_2O bands at 1.1, 1.3, and $1.9 \mu\text{m}$ are clearly evident, as are the $1.25 \mu\text{m}$ K I doublet lines. The $1.17 \mu\text{m}$ K I doublet lines are obscured by noise at the bottom of the $1.1 \mu\text{m}$ H_2O and CH_4 absorption trough. CIA H_2 is responsible for the slope on the red side of the K-band peak. These features are characteristic of mid- and late-type T dwarfs (Burgasser et al. 2002b; Geballe et al. 2002), and confirm 2MASS 1503+2525 as a bona-fide T dwarf.

4. THE SPECTRAL TYPE AND SPECTROPHOTOMETRIC DISTANCE OF 2MASS 1503+2525

We derived the spectral type of 2MASS 1503+2525 using the independent classification schemes of Burgasser et al. (2002b) and Geballe et al. (2002). Spectral indices for both schemes sample the major H_2O and CH_4 bands in T dwarf spectra, while the scheme of Burgasser et al. also includes color indices between the various spectral peaks. Measured ratio values and individual subtype determinations are listed in Table 3, following the procedures outlined by these two classification schemes. The substantial scatter amongst the Geballe et al. indices is caused by sampling low signal-to-noise regions in the Gemini Camera spectrum; similarly, the $2.2 \mu\text{m}$ CH_4 , K/J $2.11 \mu\text{m}/2.07 \mu\text{m}$ indices in the Burgasser et al. scheme are affected by the poor signal-to-noise in the K-band spectrum. Nonetheless, both yield a common spectral type of T5.5.

We confirm this spectral type in Figure 4 by comparing the J- and H-band spectra of 2MASS 1503+2525 to UKIRT CGS4 (Mountain et al. 1990) spectral data of the T5 2MASS 0559–1404 (Burgasser et al. 2000b; Geballe et al. 2002) and the T6 SDSS 1624+0029 (Strauss et al. 1999). Adequate comparison can be made between these data, as their spectral resolutions are similar ($\lambda/\Delta\lambda \sim 400$ –500) and all have been corrected for telluric absorption and flux calibrated. 2MASS 1503+2525 has stronger H_2O and CH_4 absorption bands than 2MASS 0559–1404,

but weaker bands than SDSS 1624+0029, consistent with a median type of T5.5.

As noted above, the colors of 2MASS 1503+2525 are quite blue, for its assigned spectral type of T5.5, suggestive of either a later spectral type or suppression of K-band flux, as is observed in the possibly high surface gravity or metal-poor T dwarf 2MASS 0937+2931 (Burgasser et al. 2002b). However, within the photometric uncertainties these colors are still consistent with other mid-type T dwarfs such as the T5 2MASS 0755+2212 ($J-H = 0.06 \pm 0.17$, $H-K_s = -0.08 \pm 0.26$) and the T5.5 2MASS 1534-2952AB ($J-H = 0.03 \pm 0.13$, $H-K_s = 0.03 \pm 0.16$; Burgasser et al. 2002, 2003). Higher accuracy photometry of a sample of mid-type T dwarfs is required to determine if the colors of 2MASS 1503+2525 are unusually blue for its spectral type.

Both of the comparison objects in Figure 4 have known distances from trigonometric parallax measurements (Dahn et al. 2002). Therefore, we can estimate the spectrophotometric distance of 2MASS 1503+2525 from the absolute 2MASS magnitudes of these objects and data from Table 3. If we assume this object has a spectral type and hence luminosity similar to 2MASS 0559-1404, it lies between 10.7-12.0 pc from the Sun. This estimate may be inflated if 2MASS 0559-1404 is a multiple system, as suggested by Burgasser (2001)¹⁰. On the other hand, if 2MASS 1503+2525 is a later-type, less luminous T dwarf similar to SDSS 1624+0029, it lies only 4.9-5.4 pc from the Sun. The mean spectrophotometric distance between these extremes is 8 ± 3 pc; hence, if it is single, 2MASS 1503+2525 is likely within the 10 pc RECONS distance, and is potentially a member of the 8 pc sample of Reid & Gizis (1997). Trigonometric parallax observations are currently underway for 2MASS 1503+2525 through the USNO Parallax Program (C. Dahn, priv. comm.).

Given its brightness, it is possible that 2MASS 1503+2525 could be a binary brown dwarf, as roughly 20% of late-M, L, and T dwarfs identified in magnitude-limited searches are found to have binary companions with separations $a \lesssim 10$ AU (Koerner et al. 1999; Reid et al. 2001; Close et al. 2002; Burgasser et al. 2003). The presence of a companion would increase our distance estimate by up to a factor of 1.4. Gemini images rule out a bright companion beyond $1''.2$, or 6-13 AU assuming on a distance of 5-11 pc. Because the apparent separation limit of brown dwarf binaries is similar to this observational limit (Burgasser et al. 2003), we cannot rule out the multiplicity of 2MASS 1503+2525 with any confidence; higher-resolution imaging (AO or space-based) or radial velocity monitoring are required.

5. DISCUSSION

The possibility that 2MASS 1503+2525 lies within 10 pc of the Sun raises the question of how many late-type

dwarfs are currently known in this volume. Table 4 lists all L and T dwarfs identified to date which are or may potentially be within the RECONS 10 pc horizon, based on trigonometric (Dahn et al. 2002; Perryman et al. 1997) or spectrophotometric parallax¹¹, respectively. We have used published spectrophotometric parallaxes for the L dwarfs (Cruz et al. 2003; Kirkpatrick et al. 2000) or computed them using the spectral type/absolute magnitude relations of Dahn et al. (2002), 2MASS photometry, and spectral classifications by Kirkpatrick et al. (1999). We derived distances for the T dwarfs by interpolating over spectral type (using the Burgasser et al. scheme) between the 2MASS J, H, and K_s absolute magnitudes of the L8 Gliese 584C (Kirkpatrick et al. 2000; Perryman et al. 1997), T2 SDSS 1254-0122 (Leggett et al. 2000; Dahn et al. 2002), T5 2MASS 0559-1404 (Burgasser et al. 2000b; Dahn et al. 2002), T6 SDSS 1624+0029 (Strauss et al. 1999; Dahn et al. 2002), T6.5 Gliese 229B (Leggett et al. 1999; Perryman et al. 1997, UKIRT JHK photometry), and T8 Gliese 570D (Burgasser et al. 2000a; Perryman et al. 1997). The average of spectrophotometric distances derived from 2MASS JHK_s photometry are reported.

While there are currently 313 known stars spectral type M or earlier in 227 systems within 10 pc of the Sun (T. Henry, priv. comm.), only 5 L and 10 T dwarfs meet the same criteria, if we assume that the spectrophotometric distances are accurate. In contrast, mass function simulations by Burgasser (2001), assuming conservatively $\Psi(M) \equiv dN/dM \propto M^{-0.5}$ for $0.01 M_\odot \leq M \leq 0.1 M_\odot$ ¹², predict 12 L dwarfs ($1300 \lesssim T_{eff} \lesssim 2200$ K) and 21 T dwarfs ($800 \lesssim T_{eff} \lesssim 1300$ K) in the same spatial volume, more than twice the number so far identified. The dearth of observed L and T dwarfs worsens dramatically if the substellar mass function is steeper. The primary reason for this discrepancy is quite simple - searches for these objects using large-scale surveys have not yet examined the entire sky. SDSS and DENIS programs have only searched a fraction of their final survey area, which ultimately will cover one-quarter and one-half of the sky, respectively. Searches for L and T dwarfs in the all-sky 2MASS survey have only examined roughly one-half of the sky. Hence, the expanded areal coverage and color criteria of our current search will likely identify quite a few of the missing nearby T dwarfs, particularly those with spectral types later than T4. However, some of the latest-type T dwarfs within 10 pc, and objects significantly less luminous than Gliese 570D, may be too faint to be detected by 2MASS, SDSS, or DENIS, and will remain to be uncovered in future surveys.

2MASS 1503+2525 is a key T dwarf discovery not just because of its potential proximity to the Sun, but also because it is sufficiently bright to permit both higher-resolution spectroscopic study and investigations outside the 1-2.5 μm spectral energy window. In particular, this object is a prime SIRTf target for mid-infrared observa-

¹⁰ This object was not resolved in HST observations, however, requiring the hypothetical bright companion to have a projected separation less than 0.5 AU (Burgasser et al. 2003).

¹¹ We computed spectrophotometric parallaxes for all L dwarfs in the compilations of Delfosse et al. (1997, 1999); Kirkpatrick, Beichman, & Skrutskie (1997); Kirkpatrick et al. (1999, 2000); Martín et al. (1999); Fan et al. (2000); Gizis et al. (2000); Leggett et al. (2000); Geballe et al. (2002); Gizis (2002); Hawley et al. (2002); Lodieu, Scholz, & McCaughrean (2002); Schneider et al. (2002); Cruz et al. (2003); Tinney et al. (in preparation); and Wilson et al. (in preparation); and all T dwarfs from Strauss et al. (1999); Burgasser et al. (1999, 2000b, 2002b); Leggett et al. (2000); Tsvetanov et al. (2000); and Geballe et al. (2002). See <http://spider.ipac.caltech.edu/staff/davy/ARCHIVE/index.html> for a current list of all known L and T dwarfs.

¹² Chabrier (2002) find $\Psi(M) \propto M^{-1}$, and Reid et al. (1999) find $\Psi(M) \propto M^{-1.3}$, over the same mass range.

tions, a spectral region containing diagnostics of temperature (e.g., NH_3 at $10.5\ \mu\text{m}$) and atmospheric composition (e.g., $6\text{--}9\ \mu\text{m}$ silicate bands) that is largely inaccessible due to the intrinsic faintness of these cool brown dwarfs.

We thank our referee Sandy Leggett for her useful criticisms of our manuscript and for providing the UKIRT spectral data of 2MASS 0559-1404 and SDSS 1624+0029 used in our analysis. We acknowledge useful discussions on the nearby star sample with Kelle Cruz and Todd Henry. We also thank the Gemini Camera Instrument Specialist Elinor Gates and Telescope Operator Andy Tullis at Lick for assistance in the observations, and the UCO TAC for its allocation of time for this project. Adam J. B. acknowledges support by NASA through Hubble Fellowship grant HST-HF-01137.01 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA,

under contract NAS 5-26555. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. POSS-I and POSS-II images were obtained from the Digitized Sky Survey image server maintained by the Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada. This research has made use of the NASA/IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This publication makes use of data from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center, funded by the National Aeronautics and Space Administration and the National Science Foundation. Adam J. B. dedicates this publication to the memory of Dr. Charles Kincaid Witham.

REFERENCES

- Bailer-Jones, C. A. L., & Mundt, R. 1999, *A&A*, 348, 800
 —. 2001, *A&A*, 367, 218
 Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
 Burgasser, A. J. 2001, Ph.D. Thesis, California Institute of Technology
 Burgasser, A. J., Marley, M. S., Ackerman, A. S., Saumon, D., Lodders, K., Dahn, C. C., Harris, H. C., & Kirkpatrick, J. D. 2002a, *ApJ*, 571, L151
 Burgasser, A. J., Reid, I. N., Kirkpatrick, J. D., Brown, M. E., Miskey, C. L., & Gizis, J. E. 2003, *ApJ*, submitted
 Burgasser, A. J., et al. 1999, *ApJ*, 522, L65
 —. 2000a, *ApJ*, 531, L57
 —. 2000b, *AJ*, 120, 1100
 —. 2002b, *ApJ*, 564, 421
 Chabrier, G. 2002, *ApJ*, 567, 304
 Clarke, F. J., Tinney, C. G., & Covey, K. R. 2002, *MNRAS*, 332, 361
 Close, L. M., Siegler, N., Potter, D., Brandner, W., & Liebert, J. 2002, *ApJ*, 567, L53
 Cruz, K. L., Reid, I. N., Liebert, J., Lowrance, P. J., Kirkpatrick, J. D., Gorlova, N., & Cooper, C. 2003, in *IAU Symposium 211: Brown Dwarfs*, ed. E. L. Martín (San Francisco: ASP), in preparation
 Cuby, J. G., Saracco, P., Moorwood, A. F. M., D’Odorico, S., Lidman, C., Comerón, F., & Spyromilio, J. 1999, *A&A*, 349, L41
 Dahn, C. C., et al. 2002, *AJ*, 124, 1170
 Delfosse, X., et al. 1997, *A&A*, 327, L25
 —. 1999, *A&A*, 351, L5
 Fan, X., et al. 2000, *AJ*, 119, 928
 Geballe, T. R., et al. 2002, *ApJ*, 564, 466
 Gelino, C. R., Marley, M. S., Holtzman, J. A., Ackerman, A. S., & Lodders, K. 2002, *ApJ*, 577, 433
 Gizis, J. E. 2002, *ApJ*, 575, 484
 Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. 2000, *AJ*, 120, 1085
 Golimowski, D. A., Burrows, C. S., Kulkarni, S. R., Oppenheimer, B. R., & Bruckardt, R. A. 1998, *AJ*, 115, 2579
 Guglielmo, F., Epchtein, N., Arditti, F., & Sevre, F. 1997, *A&AS*, 122, 489
 Hawley, S. L. 2002, *AJ*, 123, 3409
 Henry, T. J., Ianna, P. A., Kirkpatrick, J. D., & Jahreiβ, H. 1997, *AJ*, 114, 388
 Hess, R. 1924, in *Probleme der Astronomie: Seeliger Festschrift*, ed. H. Kienle (Springer: Berlin), p. 265
 Kirkpatrick, J. D., Beichman, C. A., & Skrutskie, M. F. 1997, *ApJ*, 476, 311
 Kirkpatrick, J. D., Reid, I. N., Liebert, J., Gizis, J. E., Burgasser, A. J., Monet, D. G., Dahn, C. C., Nelson, B., & Williams, R. J. 2000, *AJ*, 120, 447
 Kirkpatrick, J. D., et al. 1999, *ApJ*, 519, 802
 Koerner, D. W., Kirkpatrick, J. D., McElwain, M. W., & Bonaventura, N. R. 1999, *ApJ*, 526, L25
 Leggett, S. K., Toomey, D. W., Geballe, T. R., & Brown, R. H. 1999, *ApJ*, 517, L139
 Leggett, S. K., et al. 2000, *ApJ*, 536, L35
 —. 2002, *ApJ*, 564, 452
 Lodieu, N., Scholz, R.-D., & McCaughrean, M. J. 2002, *A&A*, 389, L20
 Liu, M. C., Wainscoat, R., Martín, E. L., Barris, B., & Tonry, J. 2002, *ApJ*, 568, L107
 Martín, E. L., et al. 1999, *AJ*, 118, 2466
 McLean, I. S., et al. 1993, *Proc. SPIE*, 1946, 513
 Monet, D., et al. 1998, *USNO-A2.0 Catalog* (Flagstaff: USNO)
 Mountain, C. M., Robertson, D. J., Lee, T. J., & Wade, R. 1990, *Proc. SPIE*, 1235, 25
 Nakajima, T., Oppenheimer, B. R., Kulkarni, S. R., Golimowski, D. A., Matthews, K., & Durrance, S. T. 1995, *Nature*, 378, 463
 Oppenheimer, B. R., Kulkarni, S. R., Matthews, K., & Nakajima, T. 1995, *Science*, 270, 1478
 Perryman, M. A. C., et al. 1997, *A&A*, 323, L49
 Reid, I. N., & Gizis, J. E. 1997, *AJ*, 113, 2246
 Reid, I. N., Gizis, J. E., Kirkpatrick, J. D., & Koerner, D. 2001, *AJ*, 121, 489
 Reid, I. N., et al. 1991, *PASP*, 103, 661
 —. 1999, *ApJ*, 521, 613
 Saumon, D., Bergeron, P., Lunine, J. I., Hubbard, W. B., & Burrows, A. 1994, *ApJ*, 424, 333
 Schneider, D. P., et al. 2002, *AJ*, 123, 458
 Skrutskie, M. F., et al. 1997, in *The Impact of Large-Scale Near-IR Sky Surveys*, ed. F. Garzon (Dordrecht: Kluwer), p. 25
 Strauss, M. A., et al. 1999, *ApJ*, 522, L61
 Stringov, A. R. & Sventitskii, N. S. 1968, *Tables of Spectral Lines of Neutral & Ionized Atoms* (Wilmington: IFI/Plenum Data Corp)
 Sykes, M. V., Cutri, R. M., Fowler, J. W., Tholen, D. J., Skrutskie, M. F., Price, S., & Tedesco, E. F. 2000, *Icarus*, 146, 161
 Tokunaga, A. T. 2000, in *Allen’s Astrophysical Quantities*, Fourth Edition, ed. A. N. Cox (New York: Springer-Verlag), p. 151
 Tsvetanov, Z. I., et al. 2000, *ApJ*, 531, L61
 Wilson, A. G. 1952, *Trans. I.A.U.*, 8, 335
 York, D. G., et al. 2000, *AJ*, 120, 1579
 Zapatero Osorio, M. R., Béjar, V. J. S., Martí, E. L., Rebolo, R., Barrado y Navascués, D., Mundt, R., Eisloffel, J., & Caballero, J. A. 2002, *ApJ*, 578, 536




FIG. 1.— 2MASS J-H, H-K_s Hess diagram of our T dwarf candidates, with the darkest regions indicating the highest source densities. Color constraints are indicated by dashed lines. The colors of 28 T dwarfs identified by 2MASS and SDSS are plotted as circles with error bars; arrows indicate upper limits. The colors of 2MASS 1503+2525 are distinguished by the large grey triangle. The Bessell & Brett (1988) giant and dwarf tracks are delineated by the grey lines extending redward of our sample color limits.




FIG. 2.— The field of 2MASS 1503+2525, from left to right: POSS-I (R-band), POSS-II (R-band), and 2MASS (J- and K_s -bands). Images are scaled to the same spatial resolution, $5'$ on a side, with North up and East to the left. A $10''$ box is centered on the position of 2MASS 1503+2525 in all images. The bright star $18''$ North and $103''$ East of the T dwarf is the M5 variable giant EX Boo (Guglielmo et al. 1997). Note that two bright sources below EX Boo in the 2MASS images are residual persistence sources caused by that bright star (Cutri et al. 2002).




FIG. 3.— $1.1\text{--}2.3\text{ }\mu\text{m}$ spectrum of 2MASS 1503+2525 obtained with the Lick Gemini Camera. CH $_4$, H $_2$ O, CIA H $_2$, and K I absorption features are labelled. Regions of significant telluric absorption are indicated by the $\oplus$ symbols.




FIG. 4.— Comparison of the J- (left) and H- (right) band spectra of (black lines) 2MASS 1503+2525 to (grey lines) 2MASS 0559-1404 (top) and SDSS 1624+0029 (bottom). Comparison spectral data, measured with the UKIRT CGS4 instrument, are from Geballe et al. (2002) and Strauss et al. (1999), respectively. All spectra are normalized at their flux peaks in each band.

TABLE 1
KNOWN T DWARFS IN THE 2MASS WIDE-FIELD SAMPLE.

Object ^a (1)	SpT ^b (2)	2MASS J (3)	2MASS H (4)	2MASS K _s (5)	Ref (6)
2MASS J02431371−2453298	T6	15.38±0.06	15.14±0.12	15.22±0.17	1
2MASS J04151954−0935066	T8	15.70±0.07	15.54±0.12	15.43±0.20	1
2MASS J05591914−1404488	T5	13.80±0.04	13.68±0.06	13.58±0.06	2
2MASS J07271824+1710012	T7	15.60±0.07	15.76±0.17	15.56±0.20	1
2MASS J07554795+2212169	T5:	15.73±0.07	15.67±0.15	15.75±0.21	1
SDSSp J092615.38+584720.9	T4.5	15.90±0.07	15.31±0.10	15.45±0.19	3
2MASS J09373487+2931409	T6p	14.65±0.04	14.70±0.07	15.27±0.13	1
2MASS J10475385+2124234	T6.5	15.82±0.06	15.80±0.12	> 16.4	4
2MASS J12171110−0311131	T7.5	15.86±0.07	15.75±0.13	> 15.9	4
2MASS J12255432−2739466AB	T6	15.26±0.06	15.10±0.09	15.07±0.15	4,5
2MASS J12373919+6526148 ^c	T6.5	15.81±0.07	16.28±0.24	> 15.9	4
SDSSp J134646.45−003150.4	T6	16.00±0.11	15.46±0.12	15.77±0.28	6
Gliese 570D	T8	15.32±0.06	15.27±0.09	15.24±0.16	7
2MASS J15031961+2525196	T5.5	13.94±0.03	13.86±0.04	13.96±0.06	8
2MASS J15344984−2952274AB	T5.5	14.90±0.06	14.87±0.11	14.84±0.12	1,5
2MASS J15462718−3325111	T5.5	15.63±0.06	15.45±0.10	15.49±0.18	1
2MASS J15530228+1532369	T7	15.83±0.07	15.94±0.16	15.51±0.18	1
SDSSp J162414.37+002915.6	T6	15.49±0.06	15.52±0.11	> 15.5	9
2MASS J22541892+3123498	T5	15.26±0.05	15.02±0.09	14.90±0.15	1
2MASS J23391025+1352284 ^c	T5.5	15.88±0.08	16.13±0.18	> 15.5	1
2MASS J23565477−1553111	T6	15.82±0.07	15.63±0.10	15.77±0.19	1

^aDesignations (for 2MASS discoveries) and photometry from the 2MASS PSC.

^bSpectral types from Burgasser et al. (2002b) except for SDSS 0926+5847 (Geballe et al. 2002).

^cPhotometry from another epoch satisfy our color and magnitude constraints; see Table 2.

References. — (1) Burgasser et al. (2002b); (2) Burgasser et al. (2000b); (3) Geballe et al. (2002); (4) Burgasser et al. (1999); (5) Burgasser et al. (2003); (6) Tsvetanov et al. (2000); (7) Burgasser et al. (2000a); (8) This paper; (9) Strauss et al. (1999).

TABLE 2
MULTIPLE T DWARF DETECTIONS IN THE 2MASS WPSD.

Object (1)	SpT (2)	Epoch ^a (3)	α (J2000) (4)	δ (J2000) (5)	2MASS J (6)	2MASS H (7)	2MASS K _s (8)	μ (yr ^{−1}) ^b (9)	θ (°) ^b (10)
SDSS 0926+5847	T4.5	990422	09 ^h 26 ^m 15 ^s 39	+58°47′21″19	15.60±0.08	15.62±0.17	15.41±0.24	0′′3±0′′3	220±40
		990422	09 ^h 26 ^m 15 ^s 43	+58°47′21″34	15.74±0.09	15.64±0.18	> 16.0		
		000320	09 ^h 26 ^m 15 ^s 40	+58°47′20″90	15.75±0.07	15.33±0.10	15.46±0.17		
		000320 ^c	09 ^h 26 ^m 15 ^s 38	+58°47′21″23	15.90±0.07	15.31±0.10	15.45±0.19		
2MASS 0937+2931	T6p	981124	09 ^h 37 ^m 34 ^s 81	+29°31′42″66	14.66±0.04	14.68±0.08	15.52±0.24	1′′4±0.2	150±4
		000402 ^c	09 ^h 37 ^m 34 ^s 88	+29°31′40″99	14.65±0.04	14.70±0.07	15.27±0.13		
2MASS 1237+6526	T6.5	990313	12 ^h 37 ^m 39 ^s 18	+65°26′14″59	15.81±0.07	16.28±0.24	> 15.9	— ^d	— ^d
		990313 ^c	12 ^h 37 ^m 39 ^s 20	+65°26′14″81	16.05±0.09	15.74±0.15	> 16.1		
SDSS 1346−0031	T6	990122	13 ^h 46 ^m 46 ^s 43	−00°31′50″43	15.89±0.08	16.02±0.21	> 15.7	0′′5±0′′3	275±15
		000604	13 ^h 46 ^m 46 ^s 43	−00°31′50″33	16.00±0.12	15.91±0.26	15.44±0.23		
		010203 ^c	13 ^h 46 ^m 46 ^s 34	−00°31′50″13	16.00±0.11	15.46±0.12	15.77±0.28		
		010208	13 ^h 46 ^m 46 ^s 38	−00°31′50″55	16.12±0.12	15.88±0.23	> 15.6		
Gliese 570D	T8	980516 ^c	14 ^h 57 ^m 14 ^s 96	−21°21′47″75	15.32±0.06	15.27±0.09	15.24±0.16	1′′9±0′′2	149±3
		990729	14 ^h 57 ^m 15 ^s 04	−21°21′49″72	15.08±0.06	15.35±0.14	15.30±0.21		
2MASS 1534−2952AB	T5.5	980702	15 ^h 34 ^m 49 ^s 84	−29°52′27″40	14.92±0.06	14.81±0.09	14.98±0.13	0′′31±0′′12	137±11
		980708 ^c	15 ^h 34 ^m 49 ^s 84	−29°52′27″42	14.90±0.06	14.87±0.11	14.84±0.12		
		000430	15 ^h 34 ^m 49 ^s 87	−29°52′27″82	14.97±0.07	14.72±0.07	14.85±0.12		
2MASS 2339+1352	T5.5	980929	23 ^h 39 ^m 10 ^s 21	+13°52′30″20	15.88±0.08	16.13±0.18	> 15.5	0′′83±0′′11	159±11
		001029	23 ^h 39 ^m 10 ^s 23	+13°52′28″56	16.49±0.15	15.83±0.24	> 15.6		
		001110	23 ^h 39 ^m 10 ^s 28	+13°52′28″64	16.18±0.11	15.70±0.17	> 15.8		
		001129 ^c	23 ^h 39 ^m 10 ^s 25	+13°52′28″48	16.24±0.11	15.82±0.15	16.15±0.31		

^a2MASS observation date in YYMMDD (UT).

^bProper motions derived from separate linear fits to Right Ascension and declination motion; uncertainties include the fit uncertainties and 2MASS 0′′3 positional uncertainty. Values have not been corrected for Solar motion.

^cEpoch used in 2MASS PSC.

^dEpoch difference too small to compute proper motion.

TABLE 3
T DWARF SPECTRAL CLASSIFICATION INDICES FOR 2MASS 1503+2525

Index (1)	Burgasser et al. ^a (2)	Geballe et al. ^b (3)
1.1 μ m H ₂ O	0.274 (5–6)	4.62 (6)
1.3 μ m CH ₄	0.784 (5)	—
1.5 μ m H ₂ O	0.432 (5)	3.28 (3)
1.6 μ m CH ₄	0.364 (5–6)	2.60 (6)
2.2 μ m CH ₄	0.094 (7)	10.8 (7)
H/J	0.383 (5)	—
K/J	0.131 (8)	—
2.11 μ m/2.07 μ m	1.12 (—)	—
Final Spectral Type	T5.5	T5.5

^aIndex subtypes determined by closest match to spectral standard values; final subtype is the average of the index subtypes after rejecting high and low values.

^bIndex subtypes determined from predetermined ranges; final subtype is the average of the index subtypes.

TABLE 4
KNOWN L AND T DWARFS WITHIN 10 PC OF THE SUN.

Object	SpT ^a	2MASS J	2MASS H	2MASS K _s	d (pc) ^b	Ref.
2MASS 0036+1821	L3.5	12.44±0.04	11.58±0.03	11.03±0.03	8.76±0.06	1
2MASS 0835–0819	L4.5	13.17±0.04	11.94±0.04	11.14±0.04	<i>9.1</i>	2
2MASS 1507–1627	L5	12.82±0.03	11.90±0.03	11.30±0.03	7.33±0.03	1
GJ 1001B	L5	13.10±0.03	12.05±0.02	11.40±0.03	9.55±0.10	3
DENIS 0255–4700	L8	13.23±0.03	12.19±0.02	11.53±0.03	<i>5.0</i>	4
SDSS 0423–0414	T0	14.47±0.04	13.46±0.06	12.93±0.04	<i>8.9</i>	5
2MASS 1503+2525	T5.5	13.94±0.03	13.86±0.04	13.96±0.06	<i>7.8</i>	5
2MASS 0937+2931	T6p	14.65±0.04	14.70±0.07	15.27±0.13	<i>8.8</i>	5
2MASS 0243–2453	T6	15.38±0.06	15.14±0.12	15.22±0.17	<i>9.8</i>	5
Gliese 229B	T6.5	14.32±0.05 ^c	14.35±0.05 ^c	14.42±0.05 ^c	5.77±0.04	3
2MASS 0727+1710	T7	15.60±0.07	15.76±0.17	15.56±0.20	<i>9.1</i>	5
2MASS 1553+1532	T7	15.83±0.07	15.94±0.16	15.51±0.18	<i>9.6</i>	5
2MASS 1217–0311	T7.5	15.86±0.07	15.75±0.13	> 15.9	<i>8.6</i>	5
Gliese 570D	T8	15.32±0.06	15.27±0.09	15.24±0.16	5.91±0.06	3
2MASS 0415–0935	T8	15.70±0.07	15.54±0.12	15.43±0.20	<i>6.7</i>	5

^aSpectral types from Kirkpatrick et al. (1999, 2000) and Burgasser et al. (2002b), except for 2MASS 0835–0819 (Cruz et al. 2003) and SDSS 0423–0414 (Geballe et al. 2002).

^bMeasured values are listed with uncertainties; spectrophotometric distance estimates are listed in italics.

^cUKIRT photometry from Leggett et al. (1999).

References. — (1) Dahn et al. (2002); (2) Cruz et al. (2003); (3) Perryman et al. (1997); (4) Kirkpatrick et al. (2000); (5) This paper.